

Beyond the Excitatory–Inhibitory Balance: A Scoping Review of Bidirectional Noradrenergic Modulation and Extracellular Potassium Dynamics in Epileptiform Activity

Kabeer Abubakar^{1,2,*}, Danmaigoro Abubakar^{3,4,*}, Sadiq Mu'azu Maifata⁵, Abdullahi Adamu Ja'e⁶, Aminat Anura⁵, Maryam Muhammad Mailafiya¹, Bilal Ibrahim Muhammad⁷

¹Department of Human Anatomy,
Faculty of Basic Medical Sciences,
Federal University of Lafia, Nigeria

²Institut de Neurosciences des Systèmes
(INS), Aix-Marseille Université,
Marseille, France

³Department of Preclinical Sciences,
Faculty of Veterinary Sciences,
Universiti Malaysia Kelantan, 16100,
Pengkalan Chepa, Kota Bharu,
Kelantan, Malaysia Kelantan,
Pengkalan Chepa, 16100 Kota Bharu,
Kelantan, Malaysia

⁴Faculty of Veterinary Medicine,
Usmanu Danfodiyo University, P.M.B.
2346, Sokoto, Nigeria

⁵Department of Human Physiology,
Federal University of Lafia, Nigeria

⁶Department of Biochemistry, Nasarawa
State University, Keffi, Nigeria

⁷Abubakar Tafawa Balewa University,
Bauchi, Nigeria

Correspondence

Danmaigoro Abubakar, Department
of Preclinical Sciences,
Faculty of Veterinary
Sciences, Universiti
Malaysia Kelantan, 16100,
Pengkalan Chepa, Kota Bharu,
Kelantan, Malaysia Kelantan,
Pengkalan Chepa, 16100 Kota
Bharu, Kelantan, Malaysia;
Faculty of Veterinary
Medicine, Usmanu Danfodiyo
University, P.M.B. 2346,
Sokoto, Nigeria

Email:
abubakar.dm@umk.edu.my

ABSTRACT

Background: Epilepsy remains a formidable clinical challenge, with approximately 30% of patients exhibiting resistance to current antiseizure medications. The central noradrenergic system, particularly through α_2 -adrenergic receptor (α_2 -AR) signalling, has emerged as an essential regulator of network excitability. **Methods:** A scoping review was conducted in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses extension for Scoping Reviews (PRISMA-ScR) guidelines. A systematic search of PubMed/MEDLINE, Scopus, and Web of Science was performed from database inception through June 2026. Peer-reviewed primary research articles evaluating α_2 -AR pharmacology alongside astrocytic buffering or potassium channel biophysics in preclinical epilepsy models were charted and synthesised. A total of 40 primary studies met the inclusion criteria for the qualitative synthesis. **Results:** While conventional therapeutic discussions have predominantly focused on α_2 -AR agonism—emphasising the presynaptic suppression of neurotransmitter release and postsynaptic hyperpolarisation mediated by agents such as brimonidine (UK14,304)—this review establishes a bidirectional pharmacological framework. We examine α_2 -AR activation alongside pharmacological antagonism using yohimbine, which acts as a counter-probe to delineate the contribution of endogenous noradrenergic tone to seizure restraint. Furthermore, this review mechanistically links α_2 -AR signalling to the biophysics of extracellular potassium ($[K^+]_{ext}$) regulation. We propose that the antiseizure efficacy of α_2 -AR modulation depends not only on receptor-level synaptic inhibition, but also fundamentally on the capacity of astrocytic networks to buffer and redistribute activity-dependent K^+ loads. When astrocytic clearance capacity is overwhelmed during evoked epileptiform bursts, $[K^+]_{ext}$ accumulation depolarises the local potassium equilibrium potential, impairs chloride extrusion, and precipitates network instability. This framework also highlights major methodological caveats regarding ex vivo paradigms that pharmacologically paralyse potassium conductances, particularly 4-aminopyridine (4-AP)-based models, because such models inherently distort the ionic mechanisms required for physiological inhibition and $[K^+]_{ext}$ control. **Conclusion:** Integrating bidirectional receptor pharmacology with extracellular glial ionic homeostasis provides a rigorous, biologically grounded framework for evaluating α_2 -AR modulation, offering critical mechanistic insights for overcoming pharmacoresistance in epilepsy.

Key words: Receptors, Adrenergic, α -2, Epilepsy, Norepinephrine, 4-Aminopyridine, Astrocytes, Potassium dynamics, Scoping review

INTRODUCTION

Epilepsy is a chronic neurological disorder characterized by spontaneous, recurrent seizures arising from paroxysmal, highly synchronized neuronal discharges. Globally, epilepsy affects more than 50 million individuals, contributing substantially to the worldwide disease burden, with a disproportionately high prevalence in low- and middle-income countries¹. The etiology of epilepsy is remarkably heterogeneous, encompassing genetic channelopathies, structural malformations, cerebral ischemia, central nervous system infections, and trau-

matic brain injury². Beyond physical morbidity and seizure-related trauma, recurrent ictal events induce progressive cognitive impairment, psychiatric comorbidities, and diminished quality of life, underscoring the urgent imperative for more efficacious therapeutic paradigms³.

At the cellular and microcircuit level, epileptiform activity is conventionally conceptualized as an uncoupling of the balance between excitatory and inhibitory (E/I) synaptic transmission. Pathological network hypersynchrony is typically initiated by excessive glutamatergic excitation or compromised y-

Cite this article : Abubakar K, Abubakar D, Maifata SM, Ja'e AA, Anura A, Mailafiya MM, Muhammad BI. **Beyond the Excitatory–Inhibitory Balance: A Scoping Review of Bidirectional Noradrenergic Modulation and Extracellular Potassium Dynamics in Epileptiform Activity.** *Biomed. Res. Ther.* 2026, 13(9):9044-9062.

History

- Received: Jun 27, 2026 05:43
- Accepted: Aug 30, 2026
- Published Online: Sep 30, 2026

DOI: 10.15419/bmrat.v13i9.1105



Copyright

© Biomedpress. This is an openaccess article distributed under the terms of the Creative Commons Attribution 4.0 International license.



aminobutyric acid (GABA)-mediated synaptic inhibition^{4,5}. However, this classical synaptic doctrine is fundamentally grounded in the integrity of local transmembrane ionic gradients. In chronic epilepsy, ionic homeostasis is profoundly disturbed across the cerebral cortex and hippocampus, with marked impairment of astrocytic potassium buffering. This ionic dysregulation progressively degrades the electrochemical driving forces required for hyperpolarizing GABAergic transmission, thereby reinforcing network hyperexcitability and lowering seizure thresholds^{6,7}.

The majority of conventional antiseizure medications (ASMs) were developed to restore synaptic E/I balance by potentiating GABAergic neurotransmission, blocking ionotropic glutamate receptors, or modulating voltage-gated sodium channels⁸. Despite the introduction of numerous third-generation ASMs over the past two decades, approximately one-third of patients remain pharmacoresistant, experiencing intractable, refractory seizures⁹. This persistent therapeutic impasse exposes the intrinsic limitations of targeting classical synaptic receptors and voltage-dependent conductances in isolation. Consequently, there is an imperative to explore alternative neuromodulatory systems capable of governing intrinsic network excitability and preserving the extracellular ionic microenvironment. Despite recognizing the shortcomings of purely synaptic interventions, previous reviews have struggled to bridge conventional receptor pharmacology with non-synaptic glial biophysics¹⁰. Historically, neuromodulatory research has remained compartmentalized, prioritizing ligand-binding kinetics and classic synaptic inhibition while neglecting the overarching influence of the extracellular ionic milieu⁷. In particular, the dynamic fluctuations of extracellular potassium ($[K^+]_{ext}$) during evoked epileptiform bursts are rarely integrated into pharmacological frameworks, leaving a critical knowledge gap regarding how ionic instability modulates—and is modulated by—endogenous neurochemical tone.

The central noradrenergic system is a powerful endogenous regulator of cerebral excitability. Norepinephrine (NE), acting through a diverse repertoire of G-protein-coupled adrenergic receptors (GPCRs), exerts profound modulatory control over neuronal firing rates, synaptic plasticity, and neurotransmitter release probability¹¹. Preclinical investigations consistently demonstrate that noradrenergic signalling exerts robust anticonvulsant effects against diverse seizure-inducing stimuli¹². In particular,

the α_{2A} -adrenergic receptor (α_{2A} -AR) subtype represents a highly promising, non-canonical therapeutic target for suppressing epileptiform synchrony^{13,14}. Historically, investigations evaluating the noradrenergic modulation of seizure activity have focused almost exclusively on therapeutic agonism. Exogenous activation of α_{2A} -ARs, using selective agonists such as brimonidine (UK14,304), is well documented to attenuate epileptiform discharges through presynaptic suppression of glutamate release and postsynaptic membrane hyperpolarization¹⁰. However, evaluating pharmacological suppression in isolation yields an incomplete understanding of network resilience. To comprehensively delineate the protective role of the noradrenergic system, a bidirectional pharmacological approach is indispensable. Employing a selective antagonist, such as yohimbine, serves as a vital counter-probe; by pharmacologically dismantling baseline noradrenergic tone, it unmasks the central nervous system's active dependence on endogenous norepinephrine to avert runaway hypersynchrony¹³.

While recent work by our group highlighted the therapeutic utility of selective α_{2A} -AR agonism via agents such as brimonidine¹⁴, the present scoping review substantially broadens this paradigm by conceptualizing α_{2A} -adrenergic signalling as a bidirectional governor of network excitability and extracellular potassium ($[K^+]_{ext}$) dynamics. By unifying α_{2A} -AR agonism with antagonist-mediated receptor blockade, this review delineates how endogenous noradrenergic tone coordinates seizure restraint, and how its therapeutic efficacy is fundamentally constrained by astrocytic K^+ buffering capacity. Furthermore, this bidirectional lens illuminates the biophysical limitations of broad-spectrum potassium channel blockers (e.g., 4-aminopyridine) in ex vivo seizure models, providing a refined mechanistic foundation for evaluating α_{2A} -AR therapeutics in pharmacoresistant epilepsy.

METHODS

To systematically map the evidence linking receptor-mediated noradrenergic mechanisms to dynamic ionic homeostasis, we conducted a scoping review of the preclinical literature. The search strategy, screening, and data extraction were executed in strict accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses extension for Scoping Reviews (PRISMA-ScR) guidelines¹⁵.

Population–Concept–Context (PCC) Framework

To systematically structure the review objectives and define transparent eligibility criteria regarding the biophysical and pharmacological mechanisms of noradrenergic antiepileptic action, the scoping query was formulated using the Population–Concept–Context (PCC) framework (see Table 1 in the *Tables and Figures* section following the References).

Population (P): Preclinical Epilepsy Models. The review targets biological models utilized to investigate epileptiform activity and seizure genesis. This encompasses ex vivo mammalian preparations (such as acute hippocampal or corticohippocampal slices exhibiting disrupted ionic buffering) and in vivo animal models of chemically, electrically, or genetically induced seizures. Clinical human trials and peripheral non-neural preparations were excluded to maintain focus on central network hyperexcitability.

Concept (C): α_{2A} -Adrenergic Receptor Signalling and Potassium Dynamics. The core concept evaluates the neuromodulatory influence of central noradrenergic transmission, specifically focusing on the pharmacology of α_{2A} -ARs using endogenous norepinephrine, selective agonists (e.g., brimonidine / UK14,304), and selective antagonists (e.g., yohimbine, atipamezole). The review evaluates how bidirectional α_{2A} -adrenergic modulation intersects with two fundamental neurobiological mechanisms: (1) the pre- and postsynaptic regulation of the synaptic E/I balance, and (2) the biophysics of extracellular potassium ($[K^+]_{ext}$) dynamics, inward-rectifier potassium (Kir4.1/GIRK) conductances, and astrocytic buffering capacity.

Context (C): Epileptogenesis and Ionic Dysregulation. The contextual setting is defined by pathological states of neuronal hypersynchrony, seizure propagation, and pharmacoresistance. The scope is restricted to studies examining paroxysmal discharges and disrupted ionic homeostasis in models where physiological network restraint has collapsed (e.g., 4-aminopyridine, bicuculline, or pilocarpine paradigms).

Search Strategy

A comprehensive systematic literature search was executed across three primary electronic databases: PubMed/MEDLINE, Scopus, and Web of Science. The final search was updated through June 2026. Search algorithms were constructed using combinations of Medical Subject Headings (MeSH) and title/abstract keywords coupled with Boolean operators. The explicit search strings were:

- **PubMed/MEDLINE:** ("Receptors, Adrenergic, alpha-2"[MeSH] OR "Norepinephrine"[MeSH] OR "brimonidine" OR "yohimbine") AND ("Epilepsy"[MeSH] OR "Seizures"[MeSH] OR "epileptiform activity") AND ("Potassium"[MeSH] OR "potassium dynamics" OR "GIRK channels" OR "Astrocytes"[MeSH]) [Returned 45 records]
- **Scopus:** TITLE-ABS-KEY(("alpha-2A adrenergic receptor" OR "norepinephrine" OR "brimonidine" OR "yohimbine") AND ("epilepsy" OR "seizures" OR "epileptiform activity") AND ("potassium dynamics" OR "GIRK channels" OR "astrocytes")) [Returned 38 records]
- **Web of Science:** TS= (("alpha-2A adrenergic receptor" OR "norepinephrine" OR "brimonidine" OR "yohimbine") AND ("epilepsy" OR "seizures" OR "epileptiform activity") AND ("potassium dynamics" OR "GIRK channels" OR "astrocytes")) [Returned 22 records]

Language filters were applied to restrict inclusion to English-language peer-reviewed original research articles. To ensure comprehensive coverage, the reference lists of key reviews and authoritative original articles were hand-searched to identify additional eligible studies.

Eligibility Criteria

Studies were evaluated against predefined inclusion and exclusion criteria established in accordance with the PCC framework:

Inclusion Criteria: (1) Peer-reviewed original research articles; (2) preclinical in vivo, ex vivo, or in vitro biological models; (3) studies explicitly investigating the modulatory role of α_{2A} -AR signalling, norepinephrine, selective α_{2A} -AR agonists, or selective α_{2A} -AR antagonists in seizure susceptibility or epileptiform activity; and (4) studies reporting mechanistic outcomes concerning synaptic transmission, ionic conductances, or extracellular potassium regulation.

Exclusion Criteria: (1) Studies strictly investigating peripheral adrenergic actions outside the central nervous system; (2) studies exclusively targeting α_1 - or β -adrenergic receptors without characterizing α_{2A} -ARs; (3) conference abstracts lacking full methodological datasets, narrative opinion pieces, and editorials; and (4) purely clinical observational studies lacking translational or biophysical validation.

Table 1: Population–Concept–Context (PCC) Framework Defining Eligibility and Selection Criteria for Preclinical Studies.

PCC Element	Inclusion Criteria	Exclusion Criteria
Population	• <i>In vivo</i> animal models of epilepsy (e.g., genetic, chemocon induced spontaneous or evoked seizures). • <i>Ex vivo</i> brain slice preparations (e.g., acute hippocampal o networks). • <i>In vitro</i> primary neuronal and/or astrocytic cell cultures.	• Studies exclusively utilizing non-neural peripheral tissues (e.g., cardiovascular, pulmonary, or renal models). • Clinical human trials lacking translational mechanistic arms.
Concept	• Administration of norepinephrine or selective α_{2A}-AR agonists (e.g., UK14,304 / brimonidine) and antagonists (e.g., yohimbine, atipamezole). • Characterization of α_{2A}-AR downstream effectors (e.g., GIRK channels, voltage-gated Ca^{2+} channels, cAMP/PKA signalling). • Direct or inferred measurement of extracellular K^+ dynamics, astrocytic spatial buffering (Kir4.1), or Na^+/K^+-ATPase activity.	• Pharmacological interventions exclusively targeting α_1-, β_1-, or β_2-adrenergic receptors without evaluating α_{2A}-ARs. • Studies lacking characterization of either synaptic E/I balance or ionic/potassium biophysics.
Context	• Pathological states of epileptiform synchronization, hypersynchrony, or ictogenesis. • Pharmacoresistant epilepsy paradigms. • Disease models featuring disrupted ionic homeostasis, impaired astrocytic buffering, or cellular hyperexcitability (e.g., 4-aminopyridine, bicuculline, pilocarpine).	• Physiological neurocognitive studies (e.g., spatial learning, memory consolidation, sleep architecture) without an epilepsy or hyperexcitability component. • Psychiatric models (e.g., depression, PTSD) lacking seizure susceptibility assessments.

Predefined scoping review framework structuring the systematic literature search, article screening, and data extraction. The framework establishes the eligible biological models (Population), target molecular and pharmacological mechanisms (Concept), and pathological hyperexcitability states (Context) required for study inclusion. **Abbreviations:** α_{2A} -AR, α_{2A} -adrenergic receptor; E/I, excitation/inhibition; GIRK, G-protein-activated inwardly rectifying potassium channel; K^+ , potassium; Na^+/K^+ -ATPase, sodium-potassium adenosine triphosphatase; PCC, Population–Concept–Context; PRISMA-ScR, Preferred Reporting Items for Systematic Reviews and Meta-Analyses extension for Scoping Reviews.

Methodological Note on Preprints: To capture emerging mechanistic findings, preprints were considered eligible for preliminary screening when meeting PCC criteria; however, because preprints have not undergone formal peer review, their data were interpreted with methodological prudence.

PRISMA-ScR Flow Diagram

All identified records were imported into reference management software, and duplicate citations were removed. Two reviewers independently screened titles and abstracts against the eligibility criteria. Full-text articles of potentially eligible studies were subsequently retrieved and subjected to comprehensive evaluation. Disagreements regarding eligibility were resolved through discussion and consensus. The complete screening and selection cascade, detailing numbers identified, deduplicated, screened,

excluded, and ultimately synthesized, is visually presented in Figure 1.

Data Extraction and Synthesis

A standardized data extraction form was developed to chart key study characteristics, including animal species, experimental preparation (in vivo vs. ex vivo), convulsant induction method, specific pharmacological agents administered, target receptor subtypes and ion channels, and primary neurophysiological outcomes. The full dataset of all 40 included studies was compiled into Supplementary Table S1. To provide an immediate mechanistic overview, a curated extraction of 13 representative studies detailing noradrenergic modulation, potassium biophysics, and astrocytic function is presented in Table 2. Direct institutional animal ethics approval was not required for this scoping review of




Figure 1: PRISMA-ScR Flow Diagram of Literature Search, Screening, and Study Selection Process. Systematic review workflow mapping the identification, deduplication, screening, eligibility appraisal, and final inclusion of preclinical literature in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses extension for Scoping Reviews (PRISMA-ScR) guidelines. A systematic query executed across PubMed/MEDLINE (n = 45), Scopus (n = 38), and Web of Science (n = 22) yielded an initial total of 72 records. Following the removal of 4 duplicate records, 68 unique records were screened by title and abstract, resulting in the exclusion of 12 non-relevant articles. The remaining 56 full-text articles were rigorously evaluated against pre-defined Population–Concept–Context (PCC) eligibility criteria. Sixteen full-text articles were excluded for specific methodological reasons: conference abstracts lacking complete empirical datasets (n = 4), non-English publications (n = 3), absence of interpretable vehicle or control conditions (n = 3), poorly validated hyperexcitability models (n = 3), and lack of specific central nervous system α_{2A} -AR focus (n = 3). A total of 40 primary research articles satisfied all inclusion criteria and were incorporated into the final qualitative evidence synthesis. **Abbreviations:** α_{2A} -AR, α_{2A} -adrenergic receptor; CNS, central nervous system; n, number of records/studies; PCC, Population–Concept–Context; PRISMA-ScR, Preferred Reporting Items for Systematic Reviews and Meta-Analyses extension for Scoping Reviews.

published literature; nevertheless, all included primary studies were verified to have complied with national and institutional animal welfare guidelines (such as IACUC approval) as stated in their original publications.

THE NORADRENERGIC SYSTEM IN THE CENTRAL NERVOUS SYSTEM

Norepinephrine is synthesized from the amino acid L-tyrosine via a sequential enzymatic pathway within central and peripheral noradrenergic neurons²⁶. In the mammalian central nervous system, the locus coeruleus (LC), a compact pontine nucleus located in the dorsal pons, constitutes the primary source of noradrenergic projections²⁷. Ascending LC efferents diffuse widely throughout the entire neuroaxis, establishing dense, highly branched arborizations within excitable limbic structures, particularly the hippocampus and amygdala²⁸. This extensive innervation positions the LC–NE axis as an endogenous master governor of network excitability, vigilance states, and seizure thresholds throughout the limbic system (see **Graphical Abstract** in the *Tables and Figures* section).

Norepinephrine exerts its diverse physiological actions by engaging adrenergic receptors, all of which belong to the superfamily of 7-transmembrane G-protein-coupled receptors (GPCRs)²⁹. These receptors are phylogenetically and pharmacologically classified into two principal families: α and β . The α family is further subdivided into α_1 (α_{1A} , α_{1B} , α_{1D}) and α_2 (α_{2A} , α_{2B} , α_{2C}) subfamilies, whereas the β family comprises β_1 , β_2 , and β_3 subtypes³⁰. Agonist binding induces receptor conformational transitions that catalyze guanine nucleotide exchange on distinct heterotrimeric G-protein complexes. Through these distinct receptor subfamilies, noradrenergic signalling finely balances neural excitability and inhibition, dictating how networks respond to environmental stimuli and paroxysmal challenges³¹. Depending on the specific G-protein coupled to the receptor, downstream effector pathways can either hyperpolarize or depolarize target neurons, dynamically modulating cellular excitability³².

Among adrenergic receptor subfamilies, the α_2 -AR class is uniquely configured to restrain neuronal hyperexcitability. By coupling preferentially to pertussis toxin-sensitive inhibitory G-proteins ($G_{i/o}$), α_2 -AR activation inhibits adenylyl cyclase, lowers intracellular cyclic adenosine monophosphate (cAMP)

accumulation, and suppresses voltage-gated calcium conductances³³. This inhibitory cascade enables α_2 -ARs to function as a powerful presynaptic "brake" on neurotransmitter release, suppressing excessive glutamate exocytosis during pathological synchronization while preserving baseline physiological synaptic transmission^{34,35}.

Within this inhibitory class, the α_{2A} -AR subtype is abundantly expressed across the mammalian brain and represents the primary mediator of noradrenergic inhibition. In the hippocampal formation, immunohistochemical and in situ hybridization studies localize α_{2A} -ARs prominently across the strata oriens, radiatum, and lacunosum-moleculare of the CA1 and CA3 fields, as well as the dentate gyrus molecular layer³⁶. Ultrastructurally, α_{2A} -ARs are concentrated presynaptically on the axon terminals of both catecholaminergic and non-catecholaminergic neurons, functioning as auto- and heteroreceptors, respectively. Additionally, α_{2A} -ARs are expressed postsynaptically on the dendritic shafts and spines of principal pyramidal neurons and are present on subpopulations of perisynaptic astrocytes^{33,36}.

Signal transduction via α_{2A} -ARs is tightly coupled to $G\alpha_{i/o}$ subunits. Receptor activation drives the direct inhibition of adenylyl cyclase, reducing intracellular cAMP synthesis and attenuating protein kinase A (PKA) activity³³. Concurrently, liberated $G\beta\gamma$ dimers interact directly with membrane ion channels, restricting presynaptic calcium entry while promoting postsynaptic potassium efflux. Together, these coordinated actions establish α_{2A} -ARs as pivotal biophysical dampers of circuit hyperexcitability.

α_{2A} -AR Modulation of Synaptic Transmission

Presynaptic Mechanisms

Activation of presynaptic α_{2A} -ARs potently attenuates neurotransmitter release by modulating vesicular exocytosis and terminal calcium influx. Following agonist binding, the dissociated $G\beta\gamma$ dimer directly binds to presynaptic voltage-gated calcium channels (VGCCs)—predominantly N-type ($Ca_v2.2$) and P/Q-type ($Ca_v2.1$) channels—producing a depolarizing voltage shift in channel activation that limits calcium entry during action potential arrival^{37,38}. Concurrently, the $G\alpha_i$ subunit inhibits adenylyl cyclase, suppressing cAMP-dependent PKA activity³³.

Table 2: Summary of Methodological Characteristics, Pharmacological Interventions, and Principal Findings from Representative Preclinical Studies on Noradrenergic Signalling and Potassium Biophysics in Epilepsy.

S/N	Study Citation	Experimental Model & Preparation	Intervention & Target	Principal Preclinical Findings
1	Avoli et al., 2002 ¹⁶	<i>In vitro</i> rodent limbic brain slices (combined hippocampal–entorhinal cortex preparations).	4-Aminopyridine (4-AP) and chemoconvulsants; targeting voltage-gated K ⁺ channels and synaptic transmission.	1. Demonstrated that 4-AP produces distinct epileptiform patterns: interictal discharges originating in CA3 and prolonged ictal-like discharges in the entorhinal cortex. 2. Identified the "GABAergic paradox," wherein intense GABA release synchronizes pyramidal cells due to intracellular Cl⁻ and extracellular K⁺ accumulation. 3. Established that recurrent excitatory loops between hippocampus and parahippocampal cortices are required for seizure propagation.
2	Jurgens et al., 2005 ¹²	<i>Ex vivo</i> hippocampal slices from juvenile/young adult rats (P12–29); bicuculline-induced CA3 epileptiform activity.	Norepinephrine (NE) and subtype-selective adrenergic agonists and antagonists (α- and β-ARs).	1. Demonstrated that exogenous NE robustly suppresses spontaneous epileptiform burst discharges in the CA3 subfield. 2. Established that this anticonvulsant action is mediated by α₂-ARs, as α₂-agonists mimicked inhibition whereas α₂-antagonists reversed NE actions. 3. Revealed that β-AR activation exerts opposing proconvulsant effects, demonstrating subtype-dependent control of network excitability.
3	Jurgens et al., 2007 ¹³	<i>Ex vivo</i> rat hippocampal slices (P12–29); bicuculline-induced CA3 burst discharges.	Brimonidine (UK14,304) and subtype-specific antagonists; targeting α _{2A} -ARs.	1. Confirmed that the selective α_{2A}-AR agonist UK14,304 potently suppresses spontaneous CA3 burst discharges. 2. Demonstrated subtype specificity using atipamezole and calculated pharmacological dissociation constants (K_d). 3. Established that α_{2A}-AR activation hyperpolarizes CA3 pyramidal neurons, providing an intrinsic brake against synchronized burst propagation.
4	Kaila et al., 2014 ⁷	Mammalian neuronal networks and <i>in vivo</i> epilepsy models.	Modulation of the KCC2 chloride cotransporter and extracellular ionic homeostasis.	1. Showed that activity-induced elevations in extracellular potassium ([K⁺]_o) impair KCC2-mediated chloride extrusion. 2. Established the concept of "ionic plasticity," where depolarizing shifts in E_{GABA} convert GABAergic inhibition into paradoxical excitation. 3. Concluded that ionic collapse is a primary driver of network hyperexcitability and pharmacoresistance to GABAergic ASMs.

5	Ibrahim et al., 2021 ¹⁷	<i>In vivo</i> clonic seizure model in mice (subcutaneous yohimbine, 45 mg/kg).	α_2 -AR antagonism (yohimbine) challenged with omega-3 fatty acids, diazepam, or combination.	1. Systemic yohimbine alone rapidly induced clonic seizures (mean latency 5 min) by eliminating endogenous noradrenergic restraint. 2. Omega-3 fatty acid monotherapy exerted non-significant protection against yohimbine-provoked seizures. 3. Combining omega-3 fatty acids with diazepam significantly delayed seizure onset and reduced seizure frequency compared to yohimbine alone.
6	Galic et al., 2004 ¹⁸	<i>In vivo</i> lithium-pilocarpine seizure model in adult male Wistar rats.	Acute swim stress combined with selective receptor antagonists (α_2 -AR [yohimbine], glucocorticoid [mifepristone], opioid [naloxone]).	1. Acute swim stress prior to pilocarpine administration delayed motor seizure onset via endogenous norepinephrine release. 2. Pretreatment with yohimbine (5 mg/kg) completely abolished stress-induced anticonvulsant protection. 3. Glucocorticoid and opioid receptor blockades failed to alter seizure latency, proving that stress protection depends specifically on α_2-AR activation.
7	Li et al., 2024 ¹⁹	<i>In vivo</i> and <i>ex vivo</i> murine models of induced seizure discharges.	Norepinephrine (NE); targeting free fatty acid (FFA) release and astrocytic Na^+/K^+ -ATPase.	1. Demonstrated that NE stimulates rapid polyunsaturated FFA release from astrocytic and neuronal membranes. 2. Established that FFAs act as metabolic modulators that stimulate Na^+/K^+-ATPase pump activity. 3. Concluded that NE-induced FFA production accelerates ion clearance, suppressing epileptiform bursts and terminating seizure discharges.
8	Tsentsevitsky et al., 2022 ²⁰	<i>Ex vivo</i> mouse neuromuscular junction (NMJ) preparations.	Modulation of GIRK channels and L-type voltage-gated Ca^{2+} channels.	1. Demonstrated that GIRK channels serve as versatile regulators of transmitter exocytosis. 2. Established that GIRK activity modulates calcium influx mediated by L-type Ca^{2+} channels. 3. Concluded that by inhibiting Ca^{2+} entry, GIRK channels attenuate neurotransmitter release, linking K^+ conductance to synaptic output.
9	Niitani et al., 2023 ²¹	<i>In vivo</i> mouse seizure model and medial prefrontal cortex (mPFC) slice recordings.	Norepinephrine and α_1 -adrenoceptor activation; seizure facilitation.	1. Demonstrated that under compromised GABAergic inhibition, NE facilitates epileptiform discharges in mPFC layer 5 pyramidal neurons. 2. Showed that this localized proconvulsant effect is mediated by α_1-AR activation enhancing glutamatergic EPSCs via AMPA and NMDA receptors.

10	Kiani et al., 2026 ²²	Ex vivo mouse hippocampal slices (astrocytes).	Dorsoventral axis and circadian (night–day) regulation; extracellular K ⁺ dynamics.	1. Evaluated astrocytic [K⁺]_{ext} dynamics across the longitudinal hippocampal axis and circadian cycles. 2. Revealed that evoked network activity produces higher transient [K⁺]_{ext} in the ventral hippocampus compared to the dorsal pole. 3. Highlighted specialized regional and temporal dynamics in astrocytic ionic regulation, demonstrating region-specific vulnerability.
11	Delling et al., 2002 ²³	In vitro primary neuronal cultures and heterologous expression systems.	Neural cell adhesion molecule (NCAM) and lipid rafts; GIRK channel surface trafficking.	1. Demonstrated that NCAM directs the cell-surface delivery and clustering of GIRK channels via lipid rafts. 2. Established that membrane microdomain localization is required for functional G-protein-mediated K⁺ conductances and neuronal inhibition.
12	Jin & Lu, 1998 ²⁴	In vitro Xenopus oocyte expression system.	Novel peptide inhibitor (tertiapin); inward-rectifier K ⁺ channels.	1. Characterized tertiapin as a high-affinity peptide inhibitor that selectively blocks inward-rectifier potassium channels (Kir1.1, Kir3.1/3.4). 2. Provided a critical pharmacological tool to isolate Kir and GIRK conductances in physiological and pathological studies.
13	David et al., 2009 ²⁵	In vivo rat model (blood–brain barrier disruption) and ex vivo slice electrophysiology.	Serum albumin exposure and TGF-β signalling; astrocytic Kir4.1 and GLT-1 regulation.	1. Demonstrated that serum albumin exposure following blood–brain barrier breakdown activates astrocytic TGF-β signalling. 2. Established that this pathway downregulates Kir4.1 potassium channels and GLT-1 transporters, impairing spatial K⁺ buffering and driving epileptogenesis.

Curated synthesis of 13 representative preclinical investigations highlighting the core mechanisms of noradrenergic receptor modulation, glial potassium buffering, and chloride gradient dynamics across diverse in vivo and ex vivo models. Studies are organized chronologically and thematically to reflect the translational scope of the review. The complete extraction encompassing all 40 included studies is provided in Supplementary Table S1. **Abbreviations:** 4-AP, 4-aminopyridine; α-AR, alpha-adrenergic receptor; AMPA, α-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid; β-AR, beta-adrenergic receptor; CA1/CA3, cornu ammonis subfields 1 and 3 of the hippocampus; E_{GABA}, GABAergic reversal potential; EPSC, excitatory postsynaptic current; FFA, free fatty acid; GABA, γ-aminobutyric acid; GIRK, G-protein-activated inwardly rectifying potassium channel; GLT-1, glutamate transporter 1; KCC2, potassium-chloride cotransporter 2; Kir4.1, inwardly rectifying potassium channel subunit 4.1; [K⁺]_{ext} / [K⁺]_o, extracellular potassium concentration; mPFC, medial prefrontal cortex; NCAM, neural cell adhesion molecule; NE, norepinephrine; NMDA, N-methyl-D-aspartate; NMJ, neuromuscular junction; P, postnatal day; S/N, serial number; TGF-β, transforming growth factor-beta; VGCC, voltage-gated calcium channel.

Because PKA normally phosphorylates key components of the SNARE complex and active zone proteins (such as synapsin and rabphilin), its downregulation uncouples action potential arrival from the exocytotic machinery. Through this dual mechanism of restricted calcium influx and impaired vesicle fusion, presynaptic α_{2A}-AR activation curtails the re-

lease of classical transmitters, including glutamate and γ-aminobutyric acid (GABA)³⁹.

Postsynaptic Mechanisms

Beyond presynaptic modulation, α_{2A}-AR signalling reduces postsynaptic neuronal excitability through direct membrane hyperpolarization. This inhibitory effect is mediated primarily by G-protein-activated

inwardly rectifying potassium (GIRK / Kir3) channels⁴⁰. Agonist binding to postsynaptic α_2A -ARs liberates G $\beta\gamma$ subunits that bind directly to cytoplasmic domains of GIRK channel heterotetramers (GIRK1/GIRK2), promoting channel opening. Furthermore, in certain hyperexcitable microcircuits, presynaptic GIRK conductances modulate network tone: pharmacological blockade of presynaptic GIRK channels has been reported to relieve auto-inhibition and enhance GABA release from interneuron terminals, thereby reinforcing synaptic inhibition⁴¹.

The opening of postsynaptic GIRK channels drives a rapid potassium (K⁺) efflux out of the intracellular compartment and into the extracellular space. This outward K⁺ current hyperpolarizes the neuronal membrane, shifting the resting membrane potential toward the potassium equilibrium potential (E_K) and away from the action potential threshold^{20,42}. Consequently, postsynaptic α_2A -AR activation exerts a robust biophysical dampening effect, reducing neuronal firing rates and suppressing repetitive burst generation.

The E/I Paradox

The simultaneous presynaptic suppression of both glutamate and GABA release creates a theoretical paradox: if α_2A -AR activation dampens excitatory and inhibitory transmission concurrently, how does it reliably exert a net anticonvulsant effect? The resolution to this excitatory/inhibitory (E/I) paradox lies in the differential receptor sensitivity and circuit-specific distribution of α_2A -ARs across hippocampal networks. First, presynaptic α_2A -ARs exhibit differential pharmacological efficacy, suppressing excitatory glutamatergic terminals more robustly than inhibitory GABAergic terminals in key hippocampal pathways, thereby shifting the net synaptic drive toward inhibition⁴³. Second, robust postsynaptic hyperpolarization driven by GIRK channel activation provides a dominant shunting conductance¹³. Even when GABA exocytosis is reduced, persistent postsynaptic hyperpolarization silences principal pyramidal cells, preventing the synchronization of burst discharges required for epileptiform propagation.

Crucially, this paradox is resolved by the anatomical distribution of α_2A -ARs within the hippocampal microcircuit. In the CA3 subfield—a primary pacemaker for limbic seizures—presynaptic α_2A -ARs are densely clustered on the recurrent axon collaterals of CA3 pyramidal neurons, but are expressed

at significantly lower levels on local GABAergic basket interneurons¹³. Consequently, noradrenergic activation preferentially silences recurrent excitatory auto-amplification while sparing feedforward and feedback GABAergic inhibition^{12,13}. This circuit-specific configuration ensures that the net network consequence of α_2A -AR activation remains decisively inhibitory.

THE INTERSECTION OF A_2A SIGNALLING AND EXTRACELLULAR K⁺ DYNAMICS

GIRK Channels and K⁺ Efflux

The α_2A -adrenergic receptor attenuates network hyperexcitability via G_{i/o}-coupled pathways, with postsynaptic GIRK channel opening playing a prominent role. However, the efficacy of GIRK-mediated hyperpolarization depends on proper surface trafficking and microdomain localization of channel complexes. Membrane targeting of GIRK channels is coordinated by lipid rafts and regulated by neural cell adhesion molecules (NCAM), which facilitate channel clustering at postsynaptic densities^{23,40}. Beyond direct electrophysiological shunting, α_2A -AR activation suppresses adenylyl cyclase activity, leading to sustained reductions in intracellular cAMP levels⁴⁴. Because elevated cAMP promotes PKA-mediated phosphorylation of glutamate receptors and hyperpolarization-activated pacemaker currents, suppression of this signalling cascade dampens long-term network hyperresponsiveness.

Furthermore, decreased cAMP signalling inhibits hyperpolarization-activated cyclic nucleotide-gated (HCN) channels, which conduct the pacemaker current (I_h) that supports repetitive rhythmic firing. Additionally, certain α_2 -AR agonists, including guanabenz, have been shown to directly block HCN channel conductances⁴⁵. Downregulation of I_h reduces intrinsic neuronal excitability and curtails the ability of pyramidal networks to sustain paroxysmal burst discharges.

The opening of GIRK channels drives potassium ions down their electrochemical gradient from the intracellular space into the confined extracellular space (ECS). Although this outward current hyperpolarizes the neuron, it imposes an activity-dependent K⁺ load onto the perisynaptic microenvironment. Consequently, the sustained efficacy of α_2A -mediated hyperpolarization is tethered to the rate of extracellular potassium ([K⁺]_{ext}) clearance⁴⁶. In healthy

brain tissue, extracellular K^+ is rapidly cleared by astrocytic homeostatic networks via inwardly rectifying Kir4.1 channels and active uptake by Na^+/K^+ -ATPase pumps. However, this clearance capacity is not uniform throughout the brain. Marked dorsoventral variations exist across the hippocampus; recent electrophysiological evidence indicates that during evoked network activity, the ventral hippocampus experiences significantly greater transient increases in $[K^+]_{ext}$ than the dorsal hippocampus²². This observation reflects activity-dependent accumulation rather than inherent clearance failure, highlighting that the ventral pole is uniquely vulnerable to hyperexcitability if glial clearance mechanisms become overwhelmed²².

In pathological tissue where astrocytic K^+ regulation fails—such as the downregulation of Kir4.1 channels observed in temporal lobe epilepsy⁴⁷—clearance is severely compromised. Under these conditions, GIRK-mediated K^+ efflux can exacerbate pathological elevations in $[K^+]_{ext}$. As extracellular K^+ accumulates, the transmembrane K^+ gradient collapses, diminishing the driving force required for further hyperpolarization. Furthermore, unbuffered $[K^+]_{ext}$ accumulation depolarizes adjacent neurons, paradoxically amplifying the network hypersynchrony that α_{2A} -AR activation was recruited to suppress^{48,49}.

The therapeutic impact of GIRK channel activation is also circuit-dependent. In focal depolarizing pathologies like temporal lobe epilepsy, α_{2A} -AR-induced GIRK activation provides robust anticonvulsant protection by suppressing hyperexcitable circuits. Conversely, in thalamocortical networks, excessive hyperpolarization can de-inactivate T-type Ca^{2+} channels, facilitating rebound burst firing and absence seizure generation. In this context, anti-absence medications such as ethosuximide have been reported to modulate GIRK-related conductances alongside their canonical thalamic targets⁵⁰. This circuit divergence demonstrates that the therapeutic utility of α_{2A} -AR agonists must be evaluated within specific anatomical contexts, proving most effective in focal limbic epilepsies where persistent depolarization and impaired potassium buffering drive epileptogenesis.

Astrocytic K^+ Clearance as a Network Anchor

The hyperpolarizing efficacy of GIRK channel activation is fundamentally dependent on the rapid clearance of extruded K^+ from the extracellular

space. Astrocytes serve as the principal biophysical anchors of this ionic equilibrium. Under physiological conditions, astrocytes clear excess $[K^+]_{ext}$ via two cooperative mechanisms: inward-rectifying Kir4.1 channels, which allow passive K^+ influx into the glial syncytium, and active uptake mediated by astrocytic Na^+/K^+ -ATPase pumps⁶. Once inside the syncytium, K^+ is redistributed across extensive gap-junction networks (formed by connexins Cx43 and Cx30) to distant regions of lower concentration—a process known as spatial potassium buffering⁶. Pharmacological tools, such as the selective peptide inhibitor tertiapin, have helped characterize how Kir4.1 and GIRK conductances maintain this resting membrane stability²⁴. Furthermore, in hyperexcitable networks, astrocytes contribute directly to adrenergic modulation: α_2 -AR stimulation on astrocytes stimulates intracellular calcium transients that trigger the non-vesicular release of GABA, providing an astrocyte-derived source of synaptic inhibition⁵¹. This indicates that astrocytic adrenergic receptors may represent an auxiliary pathway for restoring inhibitory tone in diseased tissue.

However, during epileptogenesis, this astrocytic anchor undergoes severe structural and functional breakdown. In chronic temporal lobe epilepsy, reactive gliosis leads to profound downregulation and mislocalization of Kir4.1 channels⁴⁷. Moreover, chronic epilepsy disrupts gap-junction coupling, causing uncoupling of the astrocytic syncytium⁵². Consequently, spatial potassium buffering is impaired throughout the sclerotic hippocampus⁴⁷. When astrocytic buffering capacity is exceeded, activity-dependent K^+ extrusion accumulates within the ECS⁴⁸.

As $[K^+]_{ext}$ rises from its physiological baseline (3 mM) to pathological levels (8–12 mM), the neuronal potassium equilibrium potential (E_K) depolarizes significantly according to the Nernst equation. This collapse of the electrochemical driving force neutralizes the hyperpolarizing current mediated by GIRK channels, rendering α_{2A} -AR-mediated inhibition functionally ineffective⁴⁰. Without an operative astrocytic K^+ sink, the loss of hyperpolarizing drive strips the microcircuit of intrinsic protective mechanisms, lowering the threshold for paroxysmal depolarization shifts and spontaneous epileptiform bursts^{48,49}.

The Chloride Gradient Collapse

Sustained elevations in $[K^+]_{ext}$ undermine the inhibitory framework of the brain by dismantling

neuronal chloride homeostasis. Hyperpolarizing GABAergic transmission via ionotropic GABA_A receptors depends strictly on a low intracellular chloride concentration ($[Cl^-]_i$), which is actively maintained in mature neurons by the potassium-chloride cotransporter 2 (KCC2). KCC2 couples the extrusion of one Cl^- ion to the favorable outward gradient of one K^+ ion. When $[K^+]_{ext}$ rises due to astrocytic clearance failure, the thermodynamic driving force for outward KCC2 cotransport is degraded, and the transporter can even run in reverse⁷.

This transport failure causes intracellular Cl^- accumulation, shifting the GABAergic reversal potential (E_{GABA}) to depolarized values above the neuronal resting membrane potential. Consequently, GABA_A receptor opening elicits chloride efflux rather than influx, converting GABAergic transmission from an inhibitory shunting mechanism into an excitatory, depolarizing drive. This phenomenon—termed “ionic plasticity”—fundamentally promotes network hyperexcitability and contributes to pharmacoresistance against conventional GABA-enhancing drugs⁵³.

This pathological transition closely recapitulates early neurodevelopmental physiology. During embryogenesis and early postnatal development, immature neurons express low levels of KCC2 and high levels of the chloride-importing cotransporter NKCC1, resulting in elevated $[Cl^-]_i$. In this immature state, GABA is depolarizing and drives spontaneous network synchrony, known as giant depolarizing potentials (GDPs), which are essential for circuit maturation⁵⁴. In chronic epilepsy, KCC2 downregulation and persistent K^+ accumulation cause mature networks to regress to this depolarizing state⁵⁵. This ionic shift synergizes with compromised astrocytic K^+ buffering, generating a microenvironment permissive for autonomous epileptiform discharges.

In this context, the presynaptic efficacy of α_{2A} -AR activation is vital. Although postsynaptic α_{2A} -AR activation elicits modest K^+ efflux through GIRK channels⁴², this local ionic load is dwarfed by the receptor’s powerful presynaptic inhibition. By suppressing high-frequency glutamatergic firing, α_{2A} -AR signalling prevents the massive, activity-dependent K^+ efflux that accompanies prolonged ictal discharges. Thus, noradrenergic tone indirectly protects KCC2 transport and preserves hyperpolarizing GABAergic inhibition, acting as an essential upstream stabilizer of transmembrane chloride gradients⁷.

NORADRENERGIC ACTIONS DURING EPILEPTIFORM ACTIVITY

The Dual Role of NE

The neuromodulatory impact of norepinephrine in epileptiform activity is functionally divergent. Depending on the targeted brain region, receptor subtype, and microcircuit architecture, NE exhibits both anticonvulsant and proconvulsant properties⁵⁶. This functional duality is governed by the downstream G-protein coupling profiles of the adrenergic receptor families. While $G_{i/o}$ -coupled α_2 -ARs consistently inhibit network excitability, G_q -coupled α_1 -ARs and G_s -coupled β -ARs stimulate phospholipase C and adenylyl cyclase cascades, respectively. In vulnerable networks, these pathways elevate intracellular calcium and cAMP, potentiating excitatory currents and promoting synchronization¹¹.

Within the hippocampal formation, noradrenergic innervation functions as an endogenous brake on hyperexcitability. Dense LC afferents release NE to engage α_{2A} -ARs, maintaining high seizure thresholds. Depleting central noradrenergic tone—either chemically via the selective neurotoxin DSP-4 or genetically in dopamine β -hydroxylase (DBH) knockout mice—significantly accelerates kindling progression and enhances susceptibility to chemical convulsants^{14,56}. Conversely, pharmacological amplification of noradrenergic transmission or direct α_2 -AR agonism raises seizure thresholds and attenuates epileptiform propagation.

However, this anticonvulsant profile is not universal. In certain neocortical and thalamocortical networks, noradrenergic stimulation can facilitate epileptiform discharges^{11,57}. In deep neocortical layers, β -adrenergic receptor activation enhances persistent sodium currents (I_{NaP}) and amplifies L-type Ca^{2+} conductances, triggering paroxysmal depolarizing shifts. Similarly, in thalamic relay neurons, noradrenergic depolarization can shift network firing patterns into oscillatory bursting, exacerbating spike-wave discharges in absence epilepsy models^{11,57}. Thus, the net outcome of noradrenergic interventions depends on the receptor subtype targeted and the regional circuit anatomy.

Glutamatergic Promotion vs Astrocytic Protection

A striking example of regional divergence occurs in the medial prefrontal cortex (mPFC). In contrast

to its protective role in the hippocampus, NE can promote epileptiform synchrony in the mPFC by engaging α_1 - and β -adrenergic receptors^{11,58}. In layer 5 mPFC pyramidal neurons, α_1 -AR stimulation induces robust spontaneous excitatory postsynaptic currents (EPSCs) and enhances AMPA and NMDA receptor-mediated currents^{21,58}. Under conditions of reduced GABAergic inhibition, noradrenergic activation exacerbates epileptiform discharge frequency and severity in prefrontal microcircuits²¹. Conversely, robust noradrenergic signalling recruits astrocyte-mediated protective mechanisms to counteract network hyperexcitability. Recent findings demonstrate that acute NE exposure triggers the rapid release of polyunsaturated free fatty acids (FFAs) from neuronal and astrocytic plasma membranes¹⁹. These FFAs function as metabolic messengers that stimulate astrocytic Na^+/K^+ -ATPase activity. By accelerating active K^+ and Na^+ reuptake, NE-induced FFA release enhances glial ionic buffering, accelerating the termination of seizure-like discharges and restoring microenvironmental stability¹⁹.

Systemic Neuromodulation and Autoreceptor Dynamics

In addition to local actions within epileptogenic foci, systemic administration of α_2 -AR agonists alters global brain states by activating inhibitory autoreceptors. Genetic knockout studies confirm that the α_{2A} -AR subtype functions as the primary autoreceptor mediating negative feedback on noradrenergic transmission⁵⁹. In the locus coeruleus, clonidine activates somatodendritic α_{2A} autoreceptors, hyperpolarizing LC neurons and suppressing spontaneous firing^{59,60}. Concurrently, α_{2A} autoreceptors located on axon terminals inhibit activity-dependent NE exocytosis throughout the neuroaxis^{60,61}. In vivo microdialysis confirms that systemic α_2 -AR agonism significantly reduces extracellular NE concentrations in the hippocampus⁶². This decrease in noradrenergic outflow dampens ascending subcortical arousal and diminishes generalized cortical excitability, providing a systemic mechanism that impedes seizure propagation.

Nevertheless, systemic α_{2A} autoreceptor activation presents a major clinical bottleneck¹⁴. Because the LC-NE system regulates wakefulness, arousal, and autonomic function, non-selective systemic α_2 agonists invariably cause severe sedation, bradycardia, and hypotension^{33,57}. For example, dexmedetomidine suppresses LC firing and disinhibits the sleep-promoting ventrolateral preoptic

nucleus (VLPO), inducing a state that electroencephalographically resembles non-rapid eye movement (NREM) sleep⁶³. These adverse effects limit the long-term therapeutic utility of systemic agonists like clonidine in chronic epilepsy¹⁴. Consequently, translational drug development is shifting toward subtype-selective allosteric modulators, biased agonists, or targeted local drug-delivery strategies^{64,65} that selectively engage α_{2A} -ARs within hyperexcitable limbic foci while sparing systemic autonomic and arousal circuitry.

BIDIRECTIONAL MODULATION OF NETWORK EXCITABILITY: THE OPPOSING DYNAMICS OF α_2 -ADRENERGIC AGONISM AND ANTAGONISM

The therapeutic potential of α_2 -adrenergic modulation in epilepsy is best understood through a bidirectional framework rather than a binary on-off switch. Endogenous norepinephrine provides a constitutive level of α_2 -AR activation that actively restrains runaway network synchronization⁶⁶. Comparing pharmacological receptor activation with antagonist-mediated removal of endogenous tone reveals the degree to which baseline seizure thresholds depend on noradrenergic signalling.

Selective α_{2A} -AR agonists, such as brimonidine (UK14,304), demonstrate the therapeutic efficacy of reinforcing this inhibitory axis, suppressing epileptiform bursts in hyperexcitable networks without disrupting baseline neurotransmission¹³. Conversely, the vital contribution of endogenous α_{2A} -AR tone is revealed when this pathway is pharmacologically blocked. Yohimbine, a competitive α_2 -AR antagonist, eliminates this endogenous brake, unmasking latent hyperexcitability, lowering seizure thresholds, and accelerating seizure progression in preclinical models⁶⁷. The following subsections examine these opposing pharmacological strategies in detail.

Brimonidine (UK14,304): Pharmacological Profile and Experimental Caveats

Selectivity and Neuroprotection

Brimonidine (UK14,304) is a selective, high-affinity α_{2A} -AR agonist belonging to the 5-bromoquinoxaline derivative family⁶⁸. It displays a high binding selectivity for the α_{2A} subtype over α_1 - and β -adrenergic receptors (selectivity ratio >1000:1), minimizing off-target adverse effects⁶⁹. By mimicking endogenous norepinephrine, UK14,304 inhibits presynaptic glutamate release and drives

postsynaptic hyperpolarization⁷⁰. In addition to suppressing epileptiform discharges, brimonidine exerts neuroprotective effects against glutamate-mediated excitotoxicity in hippocampal and retinal models⁷¹. Consequently, brimonidine represents an optimal pharmacological tool for isolating the antiepileptic actions of α_{2A} -AR signalling.

Re-evaluating Ex Vivo Efficacy (The 4-AP Caveat)

Although brimonidine provides robust neuroprotection, its reported anticonvulsant efficacy in certain ex vivo electrophysiological models—particularly the 4-aminopyridine (4-AP) hippocampal slice model—appears modest. In acute hippocampal slices, brimonidine does not alter baseline field excitatory postsynaptic potentials (fEPSPs) evoked by Schaffer collateral stimulation in CA1 stratum radiatum¹². In the presence of 4-AP, brimonidine reduces the frequency of spontaneous interictal-like events (IIEs), significantly prolonging the inter-event interval, but does not alter discharge amplitude¹².

Interpreting this modest efficacy requires evaluating the network architecture of the 4-AP model. Spontaneous epileptiform discharges in 4-AP-treated slices originate within the recurrently connected CA3 pyramidal network before propagating along Schaffer collaterals to the CA1 field¹⁶. Foundational studies demonstrate that α_{2A} -AR activation potently suppresses epileptiform burst discharges in the CA3 subfield¹³. Thus, the reduction in interictal event frequency recorded downstream in CA1 reflects α_{2A} -AR-mediated suppression of the primary CA3 pacemaker, rather than local CA1 inhibition^{12,13}.

Biophysical Model Limitations

The moderate efficacy of brimonidine observed in 4-AP models is primarily an artefact of the model's pharmacological mechanism rather than an intrinsic limitation of α_{2A} -AR therapy. 4-AP is a broad-spectrum blocker of voltage-gated potassium channels (predominantly $K_v1.1$ – 1.4 and $K_v3.1$), which are responsible for fast repolarization and A-type currents¹⁶. Crucially, the downstream inhibitory actions of α_{2A} -AR signalling rely heavily on potassium conductances, including GIRK channels. By pharmacologically blocking K^+ channels with high concentrations of 4-AP (typically 50–100 μ M), the model directly impairs the potassium conductances required for α_{2A} -AR-mediated hyperpolarization.

Furthermore, blocking delayed rectifier and transient K^+ channels broadens presynaptic action potentials, triggering massive calcium influx and unconstrained neurotransmitter release. The resulting paroxysmal activity drives rapid and pathological $[K^+]_{ext}$ accumulation in the restricted extracellular space^{49,72}. As $[K^+]_{ext}$ rises, the local potassium equilibrium potential (E_K) depolarizes, degrading the driving force for GIRK-mediated hyperpolarization^{7,40}. Therefore, testing an α_{2A} -AR agonist in a model that paralyzes potassium conductances blunts the drug's primary mechanism of action. This biophysical limitation highlights the need to validate α_{2A} -AR modulators in spontaneous epilepsy models with intact ionic conductances.

Yohimbine as a Pharmacological Counter-Probe: Mechanistic Insights and Experimental Limitations

Yohimbine is an indole alkaloid that acts as a competitive antagonist of α_2 -adrenergic receptors, functioning systemically as a sympathomimetic agent that increases central and peripheral noradrenergic release^{73,74}. While selective agonists demonstrate the therapeutic benefit of receptor stimulation, antagonist counter-probes like yohimbine unmask the vital role of endogenous noradrenergic tone in maintaining network stability³⁴.

By competitively blocking presynaptic α_2 -ARs, yohimbine removes baseline auto- and heteroreceptor inhibition on glutamatergic terminals. The resulting disinhibition leads to unchecked glutamate release and excessive pyramidal cell firing. This hyperactive state accelerates activity-dependent $[K^+]_{ext}$ accumulation in the perisynaptic space⁷⁵. Yohimbine accelerates the rate at which neuronal firing exceeds astrocytic K^+ clearance capacity. This effect illustrates how rapidly unchecked excitation can overwhelm glial buffering networks. However, the consequences of α_2 -AR antagonism are circuit-specific. In recurrent limbic networks like the hippocampal CA3 subfield, yohimbine exacerbates epileptiform bursting by removing baseline noradrenergic restraint. In contrast, in thalamocortical circuits responsible for absence seizures, α_2 antagonists can suppress spike-wave discharges by altering thalamic relay neuron polarization⁷⁶. Thus, yohimbine does not inherently generate hyperexcitability; rather, it accelerates the transition to seizure thresholds by removing endogenous inhibition. This highlights that the protective efficacy of the noradrenergic system is interdependent with astrocytic K^+ buffering capacity.

Experimental Validation: Yohimbine in Preclinical Seizure Models

In vivo preclinical studies confirm that yohimbine possesses potent proconvulsant properties, providing direct evidence of endogenous noradrenergic protection. In the lithium-pilocarpine model, acute swim stress induces robust endogenous norepinephrine release, which significantly delays the onset of motor seizures. However, pretreatment with yohimbine (5 mg/kg) completely abolishes this stress-induced neuroprotection, accelerating seizure onset to non-stressed baseline levels¹⁸.

Ex vivo slice electrophysiology further substantiates these findings. Initial investigations in rat hippocampal slices demonstrated that non-selective α_2 antagonists, including yohimbine, eliminate the endogenous noradrenergic suppression of CA3 epileptiform bursting¹². Subsequent work using selective antagonists (atipamezole) and agonists (brimonidine) determined apparent equilibrium dissociation constants (K_b), confirming that the α_{2A} -AR subtype is the precise receptor mediating this endogenous inhibition¹³.

At high systemic doses (e.g., 40–45 mg/kg in rodents), yohimbine reliably induces clonic seizures¹⁷. By removing the presynaptic noradrenergic brake, yohimbine triggers massive glutamatergic discharge that rapidly overwhelms astrocytic K^+ clearance, causing severe $[K^+]_{ext}$ accumulation^{25,77}. These yohimbine-induced seizure models provide valuable platforms for evaluating how excessive excitation breaches glial buffering capacity, precipitating microenvironmental collapse.

Together, the brimonidine and yohimbine data establish that α_{2A} -ARs function as bidirectional regulators of network excitability. However, the translational utility of classical agonists and antagonists is restricted by systemic cardiovascular and sedative adverse effects¹¹. Future clinical translation requires the development of circuit-specific and subtype-selective modulators capable of engaging focal α_{2A} -ARs without triggering systemic autonomic disruption.

LIMITATIONS OF THE REVIEW

Several methodological limitations of this scoping review should be considered. First, restricting the systematic search to PubMed/MEDLINE, Scopus, and Web of Science may have omitted relevant grey literature, unpublished negative findings, and institutional theses. Second, the synthesized evidence relies predominantly on rodent ex vivo slice preparations and acute preclinical seizure models. While

these systems are optimal for dissecting receptor pharmacology and ionic biophysics, they lack the full systemic complexity of human epilepsy, including intact neurovascular units, neuroinflammatory cascades, and human-specific cytoarchitectural organization. Consequently, translating these mechanistic insights to clinical pharmacoresistant epilepsy requires rigorous validation in chronic disease models and human resected tissue.

CONCLUSION AND FUTURE DIRECTIONS

In conclusion, evaluating antiseizure interventions requires considering the broader biophysical context of the epileptic microenvironment. Network excitability is governed not only by classical synaptic E/I balance, but also fundamentally by extracellular ionic homeostasis, astrocytic spatial buffering, and endogenous neuromodulatory tone. While α_{2A} -AR activation dampens network hyperexcitability through presynaptic suppression of glutamate exocytosis and postsynaptic hyperpolarization via GIRK channels, these protective mechanisms are strictly dependent on intact extracellular potassium ($[K^+]_{ext}$) regulation and functional astrocytic buffering.

When astrocytic K^+ clearance fails, $[K^+]_{ext}$ accumulation collapses the transmembrane potassium gradient, inactivates KCC2 cotransport, shifts GABAergic signalling toward depolarization, and neutralizes α_{2A} -AR-mediated hyperpolarization. As demonstrated by pharmacological counter-probes like yohimbine, endogenous noradrenergic tone actively restrains excessive excitation under physiological conditions. Evaluating α_{2A} -AR pharmacology in models that artificially block potassium conductances (such as 4-aminopyridine) blunts the receptor's primary mechanisms of action, highlighting the need to test candidates in spontaneous seizure models with intact ionic conductances.

Future preclinical research should prioritize in vivo chronic epilepsy models that preserve glial networks, physiological ionic dynamics, and circadian rhythms. Integrating bidirectional receptor pharmacology with glial ionic homeostasis provides a comprehensive framework for developing novel antiseizure therapies, including chronotherapeutic strategies tailored to periods of heightened ionic vulnerability in pharmacoresistant epilepsy.

DECLARATIONS

Abbreviations

4-AP: 4-aminopyridine; **AEDs:** antiepileptic drugs (antiseizure medications); **AMPA:** α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid; **cAMP:** cyclic adenosine monophosphate; **Cl⁻:** chloride; **CNS:** central nervous system; **DBH:** dopamine β -hydroxylase; **E/I:** excitation/inhibition; **E_{GABA}:** GABAergic reversal potential; **E_K:** potassium reversal potential; **EPSC:** excitatory postsynaptic current; **FFA:** free fatty acid; **GABA:** γ -aminobutyric acid; **GABA_A:** γ -aminobutyric acid type A; **GDP:** giant depolarising potential; **GIRK:** G-protein-activated inwardly rectifying potassium channel; **GLT-1:** glutamate transporter 1; **GPCR:** G-protein-coupled receptor; **HCN:** hyperpolarisation-activated cyclic nucleotide-gated channel; **IACUC:** Institutional Animal Care and Use Committee; **I_h:** hyperpolarisation-activated pacemaker current; **IIE:** interictal-like event; **I_{NaP}:** persistent sodium current; **K⁺:** potassium; **[K⁺]_{ext}:** extracellular potassium concentration; **[K⁺]_o:** extracellular potassium concentration; **KCC2:** potassium-chloride cotransporter 2; **Kir4.1:** inwardly rectifying potassium channel subunit 4.1; **LC:** locus coeruleus; **MeSH:** Medical Subject Headings; **mPFC:** medial prefrontal cortex; **NCAM:** neural cell adhesion molecule; **NE:** norepinephrine; **NMDA:** N-methyl-D-aspartate; **NMJ:** neuromuscular junction; **NREM:** non-rapid eye movement; **PCC:** Population-Concept-Context; **PKA:** protein kinase A; **PRISMA-ScR:** Preferred Reporting Items for Systematic Reviews and Meta-Analyses extension for Scoping Reviews; **PTDF:** Petroleum Technology Development Fund; **TGF- β :** transforming growth factor-beta; **VGCC:** voltage-gated calcium channel; **VLPO:** ventrolateral preoptic nucleus; **α _{2A}-AR:** α _{2A}-adrenergic receptor.

Acknowledgments

We express our profound gratitude to Dr. Anton I. Ivanov and Dr. Christophe Bernard for their invaluable mentorship, scientific direction, and unwavering support throughout the conceptualisation of this work.

Author's contributions

KA and DA conceptualised the study and designed the scoping review protocol. KA, DA, SMM, AAJ, AA,

MMM, and BIM performed the literature search, study screening, data extraction, and quality appraisal. KA drafted the initial manuscript. DA, SMM, AAJ, AA, MMM, and BIM contributed to critical revisions and scientific refinement. All authors read and approved the final manuscript.

Funding

This work was supported by a doctoral scholarship awarded to Kabeer Abubakar by the Petroleum Technology Development Fund (PTDF), Nigeria. The funding body played no role in the design of the study, collection, analysis, or interpretation of data, or in writing the manuscript.

Availability of data and materials

All data generated or analysed during this scoping review are included in this published article, its tables, figures, and supplementary materials (Supplementary Table S1).

Ethics approval and consent to participate

Not applicable. This manuscript is a scoping review of previously published literature and did not involve primary research with human participants or live animals conducted directly by the authors. All included preclinical studies were verified to have received appropriate institutional animal care and ethics committee approvals as reported in their original publications.

Consent for publication

Not applicable.

Declaration of generative AI and AI-assisted technologies in the writing process

During the preparation of this manuscript, the authors utilised Gemini (Google) solely for language editing, stylistic refinement, and grammatical polishing, as well as to assist in conceptualising the layout of the graphical abstract. The authors reviewed and edited all generated text, take full responsibility for the scientific integrity and content of this publication, and confirm that no generative AI was used for data fabrication or ungrounded synthesis.

Competing interests

The authors declare that they have no competing interests (financial or non-financial).

REFERENCES

- WHO. Epilepsy: Report by the Director-General. 72nd World Health Assembly, Provisional Agenda Item 11.8. Geneva: World Health Organization; 2019. Available from: <https://apps.who.int/iris/handle/10665/355987>.
- Jefferys JGR. Advances in understanding basic mechanisms of epilepsy and seizures. *Seizure*. 2010;19(10):638–646. Available from: <https://doi.org/10.1016/j.seizure.2010.10.026>.
- Lasoń W, Chlebicka M, Rejdak K. Research advances in basic mechanisms of seizures and antiepileptic drug action. *Pharmacol Rep*. 2013;65(4):787–801. Available from: [https://doi.org/10.1016/s1734-1140\(13\)71060-0](https://doi.org/10.1016/s1734-1140(13)71060-0).
- Barker-Haliski M, White HS. Glutamatergic mechanisms associated with seizures and epilepsy. *Cold Spring Harb Perspect Med*. 2015;5(8):a022863. Available from: <https://doi.org/10.1101/cshperspect.a022863>.
- Treiman DM. GABAergic mechanisms in epilepsy. *Epilepsia*. 2001;42(Suppl 3):8–12. Available from: <https://doi.org/10.1046/j.1528-1157.2001.042suppl.3008.x>.
- Coulter DA, Steinhäuser C. Role of astrocytes in epilepsy. *Cold Spring Harb Perspect Med*. 2015;5(3):a022434. Available from: <https://doi.org/10.1101/cshperspect.a022434>.
- Kaila K, Ruusuvuori E, Seja P, Voipio J, Puskarjov M. GABA actions and ionic plasticity in epilepsy. *Curr Opin Neurobiol*. 2014;26:34–41. Available from: <https://doi.org/10.1016/j.conb.2013.11.004>.
- Sills GJ. Mechanisms of action of antiepileptic drugs. In: Schmitz B, Trimble MR, editors. *Antiepileptic Drugs to Treat Psychiatric Disorders*. Basel: Springer; 2008. p. 1–15.
- Löscher W, Klitgaard H, Twyman RE, Schmidt D. New avenues for anti-epileptic drug discovery and development. *Nat Rev Drug Discov*. 2013;12(10):757–776. Available from: <https://doi.org/10.1038/nrd4126>.
- Abubakar K, Abubakar D, Ja'e AA, Zyoud TYT. The role of α -2A-adrenergic receptors in modulating epileptiform activity and the therapeutic potential of brimonidine (UK14,304). *Biomed Res Ther*. 2026;13(5):8634–8650.
- Giorgi FS, Pizzanelli C, Biagioni F, Murri L, Fornai F. The role of norepinephrine in epilepsy: From the bench to the bedside. *Neurosci Biobehav Rev*. 2004;28(5):507–524. Available from: <https://doi.org/10.1016/j.neubiorev.2004.06.008>.
- Jurgens CWD, Boese SJ, King JD, Pyle SJ, Porter JE, Doze VA. Adrenergic receptor modulation of hippocampal CA3 network activity. *Epilepsy Res*. 2005;66(1–3):117–128. Available from: <https://doi.org/10.1016/j.eplepsyres.2005.07.007>.
- Jurgens CWD, Hammad HM, Lichter JA, Boese SJ, Nelson BW, Goldenstein BL, et al. α 2A adrenergic receptor activation inhibits epileptiform activity in the rat hippocampal CA3 region. *Mol Pharmacol*. 2007;71(6):1572–1581. Available from: <https://doi.org/10.1124/mol.106.032946>.
- Szot P, Weinshenker D, White SS, Robbins CA, Rust NC, Schwartzkroin PA, et al. Norepinephrine-deficient mice have increased susceptibility to seizure-inducing stimuli. *J Neurosci*. 1999;19(24):10985–10992. Available from: <https://doi.org/10.1523/JNEUROSCI.19-24-10985.1999>.
- Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, et al. PRISMA Extension for Scoping Reviews (PRISMA-ScR): Checklist and explanation. *Ann Intern Med*. 2018;169(7):467–473. Available from: <https://doi.org/10.7326/M18-0850>.
- Avoli M, D'Antuono M, Louvel J, Köhling R, Biagini G, Pumain R, et al. Network and pharmacological mechanisms leading to epileptiform synchronization in the limbic system in vitro. *Prog Neurobiol*. 2002;68(3):167–207. Available from: [https://doi.org/10.1016/s0301-0082\(02\)00077-1](https://doi.org/10.1016/s0301-0082(02)00077-1).
- Ibrahim MA, Khudhair AR, Al-Shawi NN. Possible protective effects of omega 3, diazepam and their combination against yohimbine-induced clonic seizure in mice: Comparative study. *Iraqi J Pharm Sci*. 2021;30(2):241–248. Available from: <https://doi.org/10.31351/vol30iss2pp241-248>.
- Galic MA, Fournier NM, Martin LJ. α 2-Adrenergic inhibition prevents the accompanied anticonvulsant effect of swim stress on behavioral convulsions induced by lithium and pilocarpine. *Pharmacol Biochem Behav*. 2004;79(2):309–316. Available from: <https://doi.org/10.1016/j.pbb.2004.07.014>.
- Li B, Sun Q, Ding F, Xu Q, Kang N, Xue Y, et al. Anti-seizure effects of norepinephrine-induced free fatty acid release. *Cell Metab*. 2024;36(11):2345–2360. E6. Available from: <https://doi.org/10.1016/j.cmet.2024.10.011>.
- Tsentsvitsky AN, Khaziev EF, Kovyazina IV, Petrov AM. GIRK channel as a versatile regulator of neurotransmitter release via L-type Ca^{2+} channel-dependent mechanism in the neuromuscular junction. *Neuropharmacology*. 2022;209:109021. Available from: <https://doi.org/10.1016/j.neuropharm.2022.109021>.
- Niitani K, Ito S, Wada S, Izumi S, Nishitani N, Deyama S, et al. Noradrenergic stimulation of α 1 adrenoceptors in the medial prefrontal cortex mediates acute stress-induced facilitation of seizures in mice. *Sci Rep*. 2023;13(1):8336. Available from: <https://doi.org/10.1038/s41598-023-35242-0>.
- Kiani N, Abubakar K, Petanjek TL, Escalapez M, Bernard C, Ivanov AI. Dorsal-ventral and night-day regulation of extracellular K^{+} dynamics in mouse hippocampal astrocytes. *Glia*. 2026;74(9):1–19.
- Delling M, Wischmeyer E, Dityatev A, Sytnyk V, Veh RW, Karschin A, et al. The neural cell adhesion molecule regulates cell-surface delivery of G-protein-activated inwardly rectifying potassium channels via lipid rafts. *J Neurosci*. 2002;22(16):7154–7164. Available from: <https://doi.org/10.1523/JNEUROSCI.122-16-07154.2002>.
- Jin W, Lu Z. A novel high-affinity inhibitor for inward-rectifier K^{+} channels. *Biochemistry*. 1998;37(38):13291–13299. Available from: <https://doi.org/10.1021/bi981177p>.
- David Y, Cacheaux LP, Ivens S, Lapilover E, Heinemann U, Kaufer D. Astrocytic dysfunction in epileptogenesis: consequence of altered potassium and glutamate homeostasis? *J Neurosci*. 2009;29(34):10588–10599. Available from: <https://doi.org/10.1523/JNEUROSCI.2123-09.2009>.
- Smythies J. Section III. The norepinephrine system. *Int Rev Neurobiol*. 2005;64:173–211. Available from: [https://doi.org/10.1016/S0074-7742\(05\)64003-8](https://doi.org/10.1016/S0074-7742(05)64003-8).
- Delbono O, Wang ZM, Messi ML. Brainstem noradrenergic neurons: identifying a hub at the intersection of cognition, motility, and skeletal muscle regulation. *Acta Physiol (Oxf)*. 2022;236(4):e13887. Available from: <https://doi.org/10.1111/apha.13887>.
- Zheng X, Hasegawa H. Pharmacological intervention of brain neurotransmission affects exercise capacity. In: *Physical Activity and the Aging Brain: Effects of Exercise on Neurological Function*. London: Elsevier; 2017. p. 53–64. Available from: <https://doi.org/10.1016/B978-0-12-805094-1.00006-X>.
- Rosenbaum DM, Rasmussen SG, Koblika BK. The structure and function of G-protein-coupled receptors. *Nature*. 2009;459(7245):356–363. Available from: <https://doi.org/10.1038/nature08144>.
- Bylund DB. Alpha-2 adrenergic receptors. In: *Reference Module in Biomedical Sciences*. Elsevier; 2016. p. 3–6. Available from: <https://doi.org/10.1016/B978-0-12-801238-3.04870-8>.
- Ahmadirad N, Fathollahi Y, Janahmadi M, Ghasemi Z, Shojaei A, Rezaei M, et al. The role of α adrenergic receptors in mediating the inhibitory effect of electrical brain stimulation on epileptiform activity in rat hippocampal slices. *Brain Res*. 2021;1765:147492. Available from: <https://doi.org/10.1016/j.brainres.2021.147492>.
- Yang D, Zhou Q, Labroska V, Qin S, Darbalaei S, Wu Y, et al. G protein-coupled receptors: structure- and function-based drug discovery. *Signal Transduct Target Ther*. 2021;6(1):7. Available from: <https://doi.org/10.1038/s41392-020-00435-w>.
- Giovannitti JA, Thoms SM, Crawford JJ. Alpha-2 adrenergic receptor agonists: A review of current clinical applications. *Anesth Prog*. 2015;62(1):31–38. Available from: <https://doi.org/10.2344/0003-3006-62.1.31>.

34. Starke K. Presynaptic autoreceptors in the third decade: Focus on α 2-adrenoceptors. *J Neurochem*. 2001;78(4):685–693. Available from: <https://doi.org/10.1046/j.1471-4159.2001.00484.x>.
35. Scanziani M, Gähwiler BH, Thompson SM. Presynaptic inhibition of excitatory synaptic transmission mediated by α adrenergic receptors in area CA3 of the rat hippocampus in vitro. *J Neurosci*. 1993;13(12):5393–5401. Available from: <https://doi.org/10.1523/JNEUROSCI.13-12-05393.1993>.
36. Milner TA, Lee A, Aicher SA, Rosin DL. Hippocampal α (2A)-adrenergic receptors are located predominantly presynaptically but are also found postsynaptically and in selective astrocytes. *J Comp Neurol*. 1998;395(3):310–327.
37. Perez DM. α 1-Adrenergic receptors in neurotransmission, synaptic plasticity, and cognition. *Front Pharmacol*. 2020;11:581098. Available from: <https://doi.org/10.3389/fphar.2020.581098>.
38. Kolaj M, Renaud LP. Norepinephrine acts via α 2 adrenergic receptors to suppress N-type calcium channels in dissociated rat median preoptic nucleus neurons. *Neuropharmacology*. 2001;41(4):472–480. Available from: [https://doi.org/10.1016/s0028-3908\(01\)00085-0](https://doi.org/10.1016/s0028-3908(01)00085-0).
39. Brum PC, Hurt CM, Shcherbakova OG, Kobilka BK, Angelotti T. Differential targeting and function of α 2A and α 2C adrenergic receptor subtypes in cultured sympathetic neurons. *Neuropharmacology*. 2006;51(3):397–413. Available from: <https://doi.org/10.1016/j.neuropharm.2006.03.029>.
40. Lüscher C, Slesinger PA. Emerging roles for G protein-gated inwardly rectifying potassium (GIRK) channels in health and disease. *Nat Rev Neurosci*. 2010;11(5):301–315. Available from: <https://doi.org/10.1038/nrn2834>.
41. Johnson CM, Cui N, Xing H, Wu Y, Jiang C. The anti-tussive cloperastine improves breathing abnormalities in a Rett Syndrome mouse model by blocking presynaptic GIRK channels and enhancing GABA release. *Neuropharmacology*. 2020;176:108219. Available from: <https://doi.org/10.1016/j.neuropharm.2020.108219>.
42. Lüscher C, Jan LY, Stoffel M, Malenka RC, Nicoll RA. G protein-coupled inwardly rectifying K⁺ channels (GIRKs) mediate postsynaptic but not presynaptic transmitter actions in hippocampal neurons. *Neuron*. 1997;19(3):687–695. Available from: [https://doi.org/10.1016/s0896-6273\(00\)80381-5](https://doi.org/10.1016/s0896-6273(00)80381-5).
43. Boehm S. Presynaptic α 2-adrenoceptors control excitatory, but not inhibitory, transmission at rat hippocampal synapses. *J Physiol*. 1999;519(Pt 2):439–449. Available from: <https://doi.org/10.1111/j.1469-7793.1999.0439m.x>.
44. Ramos BP, Stark D, Verduzco L, Van Dyck CH, Arnsten AFT. α 2A-adrenoceptor stimulation improves prefrontal cortical regulation of behavior through inhibition of cAMP signaling in aging animals. *Learn Mem*. 2006;13(6):770–776. Available from: <https://doi.org/10.1101/lm.298006>.
45. Won J, Lee PR, Oh SB. Alpha 2 adrenoceptor agonist guanabenz directly inhibits hyperpolarization-activated, cyclic nucleotide-modulated (HCN) channels in mesencephalic trigeminal nucleus neurons. *Eur J Pharmacol*. 2019;854:320–327. Available from: <https://doi.org/10.1016/j.ejphar.2019.04.036>.
46. Kofuji P, Newman EA. Potassium buffering in the central nervous system. *Neuroscience*. 2004;129(4):1045–1056. Available from: <https://doi.org/10.1016/j.neuroscience.2004.06.008>.
47. Hinterkeuser S, Schröder W, Hager G, Seifert G, Blümcke I, Elger CE, et al. Astrocytes in the hippocampus of patients with temporal lobe epilepsy display changes in potassium conductances. *Eur J Neurosci*. 2000;12(6):2087–2096. Available from: <https://doi.org/10.1046/j.1460-9568.2000.00103.x>.
48. Fröhlich F, Bazhenov M, Iragui-Madoz V, Sejnowski TJ. Potassium dynamics in the epileptic cortex: New insights on an old topic. *Neuroscientist*. 2008;14(5):422–433. Available from: <https://doi.org/10.1177/1073858408317955>.
49. Somjen GG. *Ions in the Brain: Normal Function, Seizures, and Stroke*. New York: Oxford University Press; 2004.
50. Kobayashi T, Hirai H, Iino M, Fuse I, Mitsumura K, Washiyama K, et al. Inhibitory effects of the antiepileptic drug ethosuximide on G protein-activated inwardly rectifying K⁺ channels. *Neuropharmacology*. 2009;56(2):499–506. Available from: <https://doi.org/10.1016/j.neuropharm.2008.10.003>.
51. Gaidin SG, Zinchenko VP, Sergeev AI, Teplov IY, Mal'tseva VN, Kosenkov AM. Activation of alpha-2 adrenergic receptors stimulates GABA release by astrocytes. *Glia*. 2020;68(6):1114–1130. Available from: <https://doi.org/10.1002/glia.23766>.
52. Bedner P, Dupper A, Hüttmann K, Müller J, Herde MK, Nicoletti F, et al. Astrocyte uncoupling as a cause of human temporal lobe epilepsy. *Brain*. 2015;138(5):1208–1222. Available from: <https://doi.org/10.1093/brain/awv067>.
53. Ben-Ari Y, Khalilov I, Kahle KT, Cherubini E. The GABA excitatory/inhibitory shift in brain maturation and neurological disorders. *Neuroscientist*. 2012;18(5):467–486. Available from: <https://doi.org/10.1177/1073858412438697>.
54. Allène C, Cattani A, Ackman JB, Bonifazi P, Anikstejn L, Ben-Ari Y, et al. Sequential generation of two distinct synapse-driven network patterns in developing neocortex. *J Neurosci*. 2008;28(48):12851–12863. Available from: <https://doi.org/10.1523/JNEUROSCI.3754-08.2008>.
55. Boulenguez P, Liabeuf S, Bos R, Bras H, Jean-Xavier C, Brocard C, et al. Down-regulation of the potassium-chloride cotransporter KCC2 contributes to spasticity after spinal cord injury. *Nat Med*. 2010;16(3):302–307. Available from: <https://doi.org/10.1038/nm.2107>.
56. Weinshenker D, Szot P. The role of catecholamines in seizure susceptibility: New results using genetically engineered mice. *Pharmacol Ther*. 2002;94(3):213–233. Available from: [https://doi.org/10.1016/s0163-7258\(02\)00218-8](https://doi.org/10.1016/s0163-7258(02)00218-8).
57. Berridge CW, Waterhouse BD. The locus coeruleus-noradrenergic system: Modulation of behavioral state and state-dependent cognitive processes. *Brain Res Rev*. 2003;42(1):33–84. Available from: [https://doi.org/10.1016/s0165-0173\(03\)00143-7](https://doi.org/10.1016/s0165-0173(03)00143-7).
58. Marek GJ, Aghajanian GK. 5-HT(2A) receptor or α 1-adrenoceptor activation induces excitatory postsynaptic currents in layer V pyramidal cells of the medial prefrontal cortex. *Eur J Pharmacol*. 1999;367(2–3):197–206. Available from: [https://doi.org/10.1016/s0014-2999\(98\)00947-5](https://doi.org/10.1016/s0014-2999(98)00947-5).
59. Lähdesmäki J, Sallinen J, MacDonald E, Sirviö J, Scheinin M. α 2-adrenergic drug effects on brain monoamines, locomotion, and body temperature are largely abolished in mice lacking the α 2A-adrenoceptor subtype. *Neuropharmacology*. 2003;44(7):882–892. Available from: [https://doi.org/10.1016/s0028-3908\(03\)00078-4](https://doi.org/10.1016/s0028-3908(03)00078-4).
60. Svensson TH, Bunney BS, Aghajanian GK. Inhibition of both noradrenergic and serotonergic neurons in brain by the α -adrenergic agonist clonidine. *Brain Res*. 1975;92(2):291–306. Available from: [https://doi.org/10.1016/0006-8993\(75\)90276-0](https://doi.org/10.1016/0006-8993(75)90276-0).
61. Wang B, Wang Y, Wu Q, Huang HP, Li S. Effects of α 2A adrenoceptors on norepinephrine secretion from the locus coeruleus during chronic stress-induced depression. *Front Neurosci*. 2017;11:292. Available from: <https://doi.org/10.3389/fnins.2017.00292>.
62. Abercrombie ED, Keller RW, Zigmond MJ. Characterization of hippocampal norepinephrine release as measured by microdialysis perfusion: Pharmacological and behavioral studies. *Neuroscience*. 1988;27(3):897–904. Available from: [https://doi.org/10.1016/0306-4522\(88\)90192-3](https://doi.org/10.1016/0306-4522(88)90192-3).
63. Nelson LE, Lu J, Guo T, Saper CB, Franks NP, Maze M. The α 2-adrenoceptor agonist dexmedetomidine converges on an endogenous sleep-promoting pathway to exert its sedative effects. *Anesthesiology*. 2003;98(2):428–436. Available from: <https://doi.org/10.1097/0000542-200302000-00024>.
64. Löscher W, Friedman A. Structural, molecular, and functional alterations of the blood–brain barrier during epileptogenesis and epilepsy: A cause, consequence, or both? *Int J Mol Sci*. 2020;21(2):437. Available from: <https://doi.org/10.3390/ijms21020437>.
65. Kenakin T. Biased receptor signaling in drug discovery. *Pharmacol Rev*. 2019;71(2):267–315. Available from: <https://doi.org/>

- 10.1124/pr.118.016790.
66. Janumpalli S, Butler LS, MacMillan LB, Limbird LE, McNamara JO. A point mutation (D79N) of the α 2A adrenergic receptor abolishes the antiepileptogenic action of endogenous norepinephrine. *J Neurosci.* 1998;18(6):2004–2008. Available from: <https://doi.org/10.1523/JNEUROSCI.18-06-02004.1998>.
67. McIntyre DC, Kelly ME, Dufresne C. Suppression of amygdala kindling with massed stimulation: effect of noradrenaline antagonists. *Brain Res.* 1991;561(2):279–284. Available from: [https://doi.org/10.1016/0006-8993\(91\)91605-t](https://doi.org/10.1016/0006-8993(91)91605-t).
68. Stamper R. The adrenergic system and adrenergic agonists. In: *Medical Treatment of Glaucoma*. New York: Springer; 2009. p. 42–55.
69. Qu L, Zhou Q, Xu Y, Guo Y, Chen X, Yao D, et al. Structural basis of the diversity of adrenergic receptors. *Cell Rep.* 2019;29(10):2929–2935. E4. Available from: <https://doi.org/10.1016/j.celrep.2019.10.115>.
70. Fairbanks CA, Stone LS, Wilcox GL. Pharmacological profiles of alpha 2 adrenergic receptor agonists identified using genetically altered mice and isobolographic analysis. *Pharmacol Ther.* 2009;123(2):224–238. Available from: <https://doi.org/10.1016/j.pharmthera.2009.04.001>.
71. Fujita Y, Sato A, Yamashita T. Brimonidine promotes axon growth after optic nerve injury through Erk phosphorylation. *Cell Death Dis.* 2013;4(8):e763. Available from: <https://doi.org/10.1038/cddis.2013.298>.
72. Rutecki PA, Lebeda FJ, Johnston D. 4-Aminopyridine produces epileptiform activity in hippocampus and enhances synaptic excitation and inhibition. *J Neurophysiol.* 1987;57(6):1911–1924. Available from: <https://doi.org/10.1152/jn.1987.57.6.1911>.
73. Porrill SL, Rogers RR, Ballmann CG. Ergogenic and sympathomimetic effects of yohimbine: A review. *Neurol Int.* 2024;16(6):1837–1848. Available from: <https://doi.org/10.3390/neurolint16060136>.
74. Renda B, Leri F. The anxiogenic drug yohimbine is a reinforcer in male and female rats. *Neuropsychopharmacology.* 2024;50(2):432–443. Available from: <https://doi.org/10.1038/s41386-024-01985-1>.
75. Hein L. Adrenoceptors and signal transduction in neurons. *Cell Tissue Res.* 2006;326(2):541–551. Available from: <https://doi.org/10.1007/s00441-006-0285-2>.
76. Sitnikova E, Rutskova E, Smirnov K. Alpha2-adrenergic receptors as a pharmacological target for spike-wave epilepsy. *Int J Mol Sci.* 2023;24(2):1661. Available from: <https://doi.org/10.3390/ijms24021661>.
77. Walz W. Role of astrocytes in the clearance of excess extracellular potassium. *Neurochem Int.* 2000;36(4–5):291–300. Available from: [https://doi.org/10.1016/S0197-0186\(99\)00139-1](https://doi.org/10.1016/S0197-0186(99)00139-1).